arXiv:1006.3233v1 [math-ph] 16 Jun 2010

An $su(1,1)$ algebraic approach for the relativistic Kepler-Coulomb problem

M. Salazar-Ramírez[a], D. Martínez[b],*
R. D. Mota[c] and V. D. Granados[a]

[a] Escuela Superior de Física y Matemáticas, Instituto Politécnico Nacional, Ed. 9, Unidad Profesional Adolfo López Mateos, 07738 México D F, México.

[b] Universidad Autónoma de la Ciudad de México, Plantel Cuautepec, Av. La Corona 320, Col. Loma la Palma, Delegación Gustavo A. Madero, 07160, México D. F., México.

[c] Unidad Profesional Interdisciplinaria de Ingeniería y Tecnologías Avanzadas, IPN. Av. Instituto Politécnico Nacional 2580, Col. La Laguna Ticomán, Delegación Gustavo A. Madero, 07340 México D. F., México.

Abstract

We apply the Schrödinger factorization method to the radial second-order equation for the relativistic Kepler-Coulomb problem. From these operators we construct two sets of one-variable radial operators which are realizations for the $su(1,1)$ Lie algebra. We use this algebraic structure to obtain the energy spectrum and the supersymmetric ground state for this system.

PACS: 03.65.Pm; 03.65.Fd; 03.65.Ge; 02.20.Sv
Keywords: Relativistic hydrogen atom; Factorization methods; $su(1,1)$ Lie algebra

---

* *E-mail address:* dmartinezs77@yahoo.com.mx

1 Introduction

Compact and non-compact symmetries play a central role to study many properties of quantum systems because they form the basis for selection rules that forbid the existence of certain states and processes. Also, from a given solution, symmetries allow to obtain new solutions of the dynamical equations. Moreover, the conserved quantity associated with a given symmetry represents an integrability condition.

Generators of compact and non-compact algebras for a given Hamiltonian have not been constructed by a systematic method but intuitively found and forced to close an algebra [1, 2, 3]. In particular, the $su(1,1) \sim so(2,1)$ non-compact algebra is the smallest dynamical symmetry admitting infinite-dimensional representations. This symmetry has been successfully applied to formulate algebraic approaches to many non-relativistic quantum problems [4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14], where the corresponding realizations were given in terms of two variables. Also, some one-variable realizations for the $su(1,1)$ algebra have been introduced [1]. Recently, it has been shown that the Schrödinger factorization operators can be used to construct the $su(1,1)$ algebra generators for two- [15] and N-dimensional systems [16].

The relativistic Kepler-Coulomb problem is one of the few exactly solvable potentials in physics and it has been studied in several ways: analytical [17, 18, 19, 20, 21, 22], factorization (algebraic) methods [23, 24, 25], shape-invariance [26] and SUSY QM for the first- [18, 27] and second-order equations [28]. The solubility of this problem is due to the conservation of the total angular momentum, the Dirac and Lippmann-Johnson operators [29]. The first two ones are due to the existence of spin, and the later explains the degeneracy in the eigenvalues of the Dirac operator of the energy spectrum and it reduces to the Runge-Lenz vector in the non-relativistic limit. Symmetries and SUSY QM are not independent. Indeed, it has been shown that supersymmetry is generated by the Lippmann-Johnson operator [29].

In a series of papers some realizations of compact and non-compact Lie algebras for the uncoupled second-order radial equations corresponding to the relativistic Kepler-Coulomb problem have been introduced [30, 31, 32]. However, in all these works the origin of the generators was not given. Moreover, in order to close the $su(1,1)$ Lie algebra these generators were forced to depend on an extra variable which plays the role of a phase.

The aim of this paper is to study the relativistic Kepler-Coulomb problem from an $su(1,1)$ algebraic approach by using the Schrödinger factorization and without the introduction of any additional variable.

In section 2, we obtain the uncoupled second-order differential equations satisfied by the radial components. In section 3, we obtain the $su(1,1)$ algebra generators by applying the Schrödinger factorization to an uncoupled second-order differential equation. These results allow us to obtain the energy spectrum for this system. In section 4, the SUSY ground state and the action of the $su(1,1)$ algebra generators on the radial eigenstates are found. Finally, we give some concluding remarks.

2 The relativistic Kepler-Coulomb radial equation

The Dirac radial equation for the Kepler-Coulomb problem is [17, 27]

$$\begin{pmatrix} \frac{dG_k^{(1)}}{dr} \\ \frac{dG_k^{(2)}}{dr} \end{pmatrix} + \frac{1}{r}\begin{pmatrix} k & -\gamma \\ \gamma & -k \end{pmatrix}\begin{pmatrix} G_k^{(1)} \\ G_k^{(2)} \end{pmatrix} = \begin{pmatrix} 0 & \alpha_1 \\ \alpha_2 & 0 \end{pmatrix}\begin{pmatrix} G_k^{(1)} \\ G_k^{(2)} \end{pmatrix}, \tag{1}$$

where k is the eigenvalue of the Dirac operator $K = -\left(\vec{\sigma}\cdot\vec{L} + \mathbf{1}\right), \gamma = \frac{ze^2}{c\hbar}, \alpha_1 = m + E, \alpha_2 = m - E, |k| = j + \frac{1}{2}, k = \pm1, \pm2, \pm3, ..., \vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ are the Pauli matrices, $\vec{L}$ is the angular momentum operator and $\mathbf{1}$ is the 2×2 unit matrix.

By introducing the new variable $\rho = Er$ equation (1) leads to the pair of equations

$$\left(\frac{k}{s} + \frac{m}{E}\right)F_k^{(2)} = \left(\frac{d}{d\rho} + \frac{s}{\rho} - \frac{\gamma}{s}\right)F_k^{(1)}, \tag{2}$$

$$\left(\frac{k}{s} - \frac{m}{E}\right)F_k^{(1)} = \left(-\frac{d}{d\rho} + \frac{s}{\rho} - \frac{\gamma}{s}\right)F_k^{(2)}, \tag{3}$$

where

$$\begin{pmatrix} F_k^{(1)} \\ F_k^{(2)} \end{pmatrix} = D\begin{pmatrix} G_k^{(1)} \\ G_k^{(2)} \end{pmatrix} \equiv \begin{pmatrix} k+s & -\gamma \\ -\gamma & k+s \end{pmatrix}\begin{pmatrix} G_k^{(1)} \\ G_k^{(2)} \end{pmatrix} \tag{4}$$

and

$$s = \sqrt{k^2 - \gamma^2}. \tag{5}$$

From these relations we obtain the uncoupled second-order differential equations

$$\left(-\frac{d^2}{d\rho^2}+\frac{s\,(s\pm 1)}{\rho^2}+\frac{\gamma^2}{s^2}-\frac{2\gamma}{\rho}\right)F_k^{(1,2)}=\left(\frac{k}{s}+\frac{m}{E}\right)\left(\frac{k}{s}-\frac{m}{E}\right)F_k^{(1,2)}, \tag{6}$$

where the superscript $(1,2)$ corresponds to $(+,-)$ in the centrifugal term, respectively. Equation (5) allows us to rewrite equation (6) as

$$\left(-\rho^2\frac{d^2}{d\rho^2}+\xi^2\rho^2-2\gamma\rho\right)F_k^{(1)}=-s\,(s+1)\,F_k^{(1)}, \tag{7}$$

$$\left(-\rho^2\frac{d^2}{d\rho^2}+\xi^2\rho^2-2\gamma\rho\right)F_k^{(2)}=-s\,(s-1)\,F_k^{(2)}, \tag{8}$$

where

$$\xi^2=\frac{m^2}{E^2}-1. \tag{9}$$

It must be emphasized that equation (7) is formally obtained from equation (8) by making $s\to s+1$. Hence, by defining $\psi_s\equiv F_k^{(2)}$, we conclude that

$$F_k^{(1)}\propto\psi_{s+1}. \tag{10}$$

Thus, the solution for the Dirac equation in spinorial form is

$$\Phi_k\equiv\begin{pmatrix}F_k^{(1)}\\ F_k^{(2)}\end{pmatrix}=\begin{pmatrix}\psi_{s+1}\\ \psi_s\end{pmatrix}. \tag{11}$$

3 The Schrödinger Factorization and the $su(1,1)$ Lie algebra

In order to factorize the left-hand side operator of equations (8), we apply the Schrödinger factorization [33, 34]. Thus, we propose a pair of first-order differential operators such that

$$\left(\rho\frac{d}{d\rho}+a\rho+b\right)\left(-\rho\frac{d}{d\rho}+c\rho+f\right)\psi_s=g\psi_s, \tag{12}$$

where a, b, c, f and g are constants to be determined. Expanding this expression and comparing it with equation (8) we obtain

$$a=c=\pm\xi,\quad f=1+b=\mp\frac{\gamma}{\xi},\quad g=b(b+1)-s(s-1). \tag{13}$$

Using these results, equation (8) is equivalent to

$$\left(B_{\mp}\mp 1\right)B_{\pm}\psi_s=\left[\left(\frac{\gamma}{\xi}\pm\frac{1}{2}\right)^2-\left(s-\frac{1}{2}\right)^2\right]\psi_s, \tag{14}$$

where

$$B_{\pm}=\mp\rho\frac{d}{d\rho}+\xi\rho-\frac{\gamma}{\xi}. \tag{15}$$

From equation (8) we obtain the operator

$$\Sigma_3\psi_s\equiv\frac{1}{2\xi}\left(-\rho\frac{d^2}{d\rho^2}+\xi^2\rho+\frac{s\left(s-1\right)}{\rho}\right)\psi_s=\frac{\gamma}{\xi}\psi_s. \tag{16}$$

Therefore, using equation (15) we define a new pair of operators

$$\Sigma_{\pm}\equiv\mp\rho\frac{d}{d\rho}+\xi\rho-\Sigma_3. \tag{17}$$

In order to establish the properties of the operators Σ_3 and $\Sigma_{\pm}$ we define the inner product on the Hilbert space spanned by the radial eigenfunctions of the relativistic Kepler-Coulomb problem as [35]

$$\left(\phi,\zeta\right)\equiv\int_0^{\infty}\phi^*\left(\rho\right)\zeta\left(\rho\right)\rho^{-1}d\rho. \tag{18}$$

Thus, it follows that the operator Σ_3 is hermitian. Moreover, using equations (17) and (18) we prove that the operators $\Sigma_{\pm}$ are hermitian conjugates,

$$\Sigma_{\pm}=\Sigma_{\mp}^{\dagger}. \tag{19}$$

By direct calculation it is immediate to show that the operators $\Sigma_{\pm}$ and Σ_3 close the $su(1,1)$ Lie algebra

$$\left[\Sigma_{\pm},\Sigma_3\right]=\mp\Sigma_{\pm}, \tag{20}$$

$$\left[\Sigma_{+},\Sigma_{-}\right]=-2\Sigma_3, \tag{21}$$

with quadratic Casimir operator

$$\Sigma^2=-\Sigma_{+}\Sigma_{-}+\Sigma_3^2-\Sigma_3. \tag{22}$$

From equations (16) and (17) we find that the eigenvalue equation for this operator is

$$\Sigma^2\psi_s = s\,(s-1)\,\psi_s. \tag{23}$$

The theory of unitary irreducible representations of the $su(1,1)$ Lie algebra has been studied in several works [35, 36] and it is based on the equations

$$C^2|\mu\,\nu\rangle = \mu(\mu+1)|\mu\,\nu\rangle, \tag{24}$$

$$C_3|\mu\,\nu\rangle = \nu|\mu\,\nu\rangle, \tag{25}$$

$$C_\pm|\mu\,\nu\rangle = \left[(\nu\mp\mu)(\nu\pm\mu\pm1)\right]^{1/2}|\mu\,\nu\pm1\rangle, \tag{26}$$

where C^2 is the quadratic Casimir operator, $\nu=\mu+q+1$, $q=0,1,2,...$ and $\mu>-1$. The last relation means that $C_+(C_-)$ are the raising (lowering) operators for ν.

Therefore, from equations (23) and (24), and (16) and (25), we find

$$\mu_s = s-1, \tag{27}$$

$$\nu_s = n_s+s=\frac{\gamma}{\xi}, \tag{28}$$

respectively, with $n_s=0,1,2,....$

If we consider that the operators $\Sigma_\pm$ leave fixed the quantum number μ_s then, from equation (27), the values of s remain fixed. This is because the change $\nu_s\to\nu_s\pm1$ induced by the operators $\Sigma_\pm$ on the basis vectors $|\mu_s\,\nu_s\rangle$ implies $n_s\to n_s\pm1$. Thus, by setting $|\mu_s\,\nu_s\rangle\to\psi_s^{n_s}$ and from equations (26), (27) and (28) we find

$$\Sigma_\pm\psi_s^{n_s} = Q_\pm^{(n_s,s)}\psi_s^{n_s\pm1}, \tag{29}$$

with $Q_\pm^{(n_s,s)} = \left[(n_s+s+1\mp s)\,(n_s+s+1\pm s\pm1)\right]^{1/2}$ a real number.

Using equations (9) and (28) we find that the energy spectrum for the lower component of the spinor given in equation (11), $\psi_s^{n_s}$, is

$$E_s = m\left[1+\frac{\gamma^2}{(n_s+s)^2}\right]^{-1/2}. \tag{30}$$

As was emphasized above, by performing the change $s\to s+1$ to $\psi_s^{n_s}$ we obtain the upper component of the spinor Φ_k, $\psi_{s+1}^{n_{s+1}}$. In this way, its corresponding differential operators are

$$\Xi_3 \equiv \frac{1}{2\xi}\left(-\rho\frac{d^2}{d\rho^2} + \xi^2\rho + \frac{s\,(s+1)}{\rho}\right), \tag{31}$$

$$\Xi_\pm \equiv \mp\rho\frac{d}{d\rho} + \xi\rho - \Xi_3 \tag{32}$$

and by direct calculation we show they satisfy the $su(1,1)$ Lie algebra

$$[\Xi_\pm, \Xi_3] = \mp\Xi_\pm, \tag{33}$$

$$[\Xi_+, \Xi_-] = -2\Xi_3. \tag{34}$$

From equation (29) we obtain that the action of the ladder operators $\Xi_\pm$ on the functions $\psi_{s+1}^{n_{s+1}}$ is

$$\Xi_\pm\psi_{s+1}^{n_{s+1}} = Q_\pm^{(n_{s+1},s+1)}\psi_{s+1}^{n_{s+1}\pm 1}. \tag{35}$$

Hence, from equation (30) we find that the energy spectrum for the function $\psi_{s+1}^{n_{s+1}}$ is

$$E_{s+1} = m\left[1 + \frac{\gamma^2}{(n_{s+1}+s+1)^2}\right]^{-1/2}. \tag{36}$$

Since $\psi_{s+1}^{n_{s+1}}$ and $\psi_s^{n_s}$ are the components for the spinor Φ_k, they must have the same energy. This means that $E_s = E_{s+1}$. Therefore, from equations (30) and (36) we obtain

$$n \equiv n_s = n_{s+1} + 1, \tag{37}$$

where $n = 0, 1, 2, 3, ...$ is the radial quantum number. Thus, the energy spectrum for the relativistic Kepler-Coulomb problem is

$$E = m\left[1 + \frac{\gamma^2}{(n+s)^2}\right]^{-1/2}, \tag{38}$$

where the positiveness of E ensures that the components of the spinor given in (11) are quadratically integrable [30]. In this way, Φ_k^n is given by

$$\Phi_k^n = \begin{pmatrix} F_{n\,k}^{(1)} \\ F_{n\,k}^{(2)} \end{pmatrix} = \begin{pmatrix} \psi_{s+1}^{n-1} \\ \psi_s^n \end{pmatrix}. \tag{39}$$

This result, which has been obtained from the theory of unitary representations, can be deduced from an analytical approach as it is shown in the next section.

4 Schrödinger and SUSY QM ground states

For $n = 0$ and from equations (29) and (37) we find that only the state

$$\psi_s^0 = N\rho^s e^{-\gamma\rho/s}, \tag{40}$$

is normalizable respect to the inner product defined in expression (18) and satisfies the differential equation $\Sigma_- \psi_s^0 = 0$. Also, for $n = 0$ and from equation (37) we find $n_{s+1} = -1$. For these values, $Q_-^{(-1,s+1)}$ results to be a complex number. Thus, from the theory of unitary representations and equation (35), the function ψ_{s+1}^{-1} is non-normalizable [36]. Since this function is not a physically acceptable solution, the spinor corresponding to $n = 0$ is

$$\Phi_k^0 = \begin{pmatrix} 0 \\ \rho^s e^{-\gamma\rho/s} \end{pmatrix}, \tag{41}$$

whom we denote as the Schrödinger ground state. Notice that Φ_k^0 is equal to the SUSY ground state for the relativistic Kepler-Coulomb problem found in [27, 29].

From an analytical approach, to determine Φ_k^n for higher levels we solve the differential equation (8) by proposing

$$F_k^{(2)} = \rho^s e^{-\xi\rho} f(\rho). \tag{42}$$

Thus, $f(\rho)$ must satisfy

$$\left[y\frac{d^2}{dy^2} + (2s - y)\frac{d}{dy} + \frac{\gamma}{\xi} - s \right] f\left(y/2\xi\right) = 0, \tag{43}$$

where $y = 2\xi\rho$. The solutions for this equation is the confluent hypergeometric function ${}_1F_1\left(-n, 2s; y\right)$. Thus, we get

$$F_{n\,k}^{(2)} = N_2 \rho^s e^{-\xi\rho}\, {}_1F_1\left(-n, 2s; 2\xi\rho\right). \tag{44}$$

In a similar way, the solution for equation (7) is

$$F_{n\,k}^{(1)} = N_1 \rho^{s+1} e^{-\xi\rho}\, {}_1F_1\left(-n + 1, 2s + 2; 2\xi\rho\right). \tag{45}$$

This equations are in agreement with our results obtained from the theory of unitary representations, equation (39). Moreover, it is known that ${}_1F_1(0, b; z) = 1$, whereas ${}_1F_1(a, b; z)$ diverges for $a > 0$. Thus, for $n = 0$

$$F_{0\,k}^{(2)} = N_s \rho^s e^{-\xi\rho}, \tag{46}$$


$F^{(2)}_{n\,1}$ $F^{(1)}_{n\,1}$
$F^{(2)}_{n\,2}$ $F^{(1)}_{n\,2}$
$F^{(2)}_{n\,3}$ $F^{(1)}_{n\,3}$
$F^{(2)}_{n\,4}$ $F^{(1)}_{n\,4}$
$F^{(2)}_{n\,5}$ $F^{(1)}_{n\,5}$
$N=5$
$N=4$
$N=3$
$N=2$
$N=1$
Σ_+
Ξ_+
Σ_-
Ξ_-
A^-
A^+
A^-
Σ_-
k
-1 1
-2 2
-3 3
-4 4
-5 5

Figure 1: Energy levels for the relativistic Kepler-Coulomb problem. The dashed lines correspond to the null component of the Schrödinger ground state and the radial quantum number $n = N - |k|$, with $N = 1, 2, 3, ...$ the principal quantum number. The action of the operators $A^{\pm}$, $\Sigma_{\pm}$ and $\Xi_{\pm}$ on the radial states is shown.

while the function $F^{(1)}_{0\,k}$ is not square-integrable and it must be taken as the zero funtion. This result is in accordance with equation (41) which has been obtained from an algebraic approach.

From the definition (37), equations (29) and (35) imply

$$\Sigma_{\pm}\psi^{n}_{s} \propto \psi^{n\pm1}_{s}, \tag{47}$$

$$\Xi_{\pm}\psi^{n-1}_{s+1} \propto \psi^{n-1\pm1}_{s+1}. \tag{48}$$

It must be notice that equation (48) is valid for any value of the radial quantum number except for $n = 0$. This is because the upper component of the spinor for $n = 1$ can not be obtained from the action of the operator Ξ_+ on the upper component of the Schrödinger ground state (equation (41)). These results are shown in Figure 1. We illustrate the action of the SUSY operators $A^{\pm}$ on the eigenstates of the relativistic Kepler-Coulomb problem [27]. Also, it is shown that the lower components for the states corresponding to $n = 0$ are annihilated by the operators Σ_- and A^-. This result and the fact that the upper component of the SUSY [27] and the Schrödinger ground states are zero imply the equality of these states.

Equations (16), (47) and (48) allow us to show that the action of the Schrödinger operators on the states ψ_s^n is

$$B_{\pm}\psi_s^n \propto \psi_s^{n\pm 1}. \tag{49}$$

A similar result can be obtained from the equation (37) and the operators $\Xi_{\pm}$. These results imply that the action of the Schrödinger operators on the components of the spinor Φ_k^n is to change only the radial quantum number n leaving fixed the Dirac quantum number $k = k(s)$.

5 Concluding Remarks

We have shown that the algebraic treatment for the radial equations of the relativistic Kepler-Coulomb problem is reduced to find the non-compact symmetries for equation (7) or (8). By applying the Schrödinger factorization we found the one-variable generators for the $su(1,1)$ Lie algebra. From the theory of unitary representations and the relation between the components of the spinor Φ_k^n we found the energy spectrum for this system in a purely algebraic way. Moreover, equations (47) and (48) imply that the $su(1,1)$ algebra generators $\Xi_{\pm}$, Ξ_3, and $\Sigma_{\pm}$, Σ_3, are represented by infinite-dimensional Hilbert subspaces of the radial quantum states. The non-compact nature of the $su(1,1)$ algebra for this problem reflects that for a fixed Dirac quantum number, the radial quantum number is bounded from below and unbounded from above. We showed that the Schrödinger ground state, which corresponds to the lowest value of the radial quantum number $n = 0$, is equal to the SUSY ground state [27, 29]. Finally, we emphasize that our treatment does not introduce an extra variable and shows the origin of the $su(1,1)$ Lie algebra generators.

Acknowledgments

This work was partially supported by SNI-México, CONACYT grant number J1-60621-I, COFAA-IPN, EDI-IPN, SIP-IPN projects numbers 20100897 and 20100518, and ADI-UACM project number 7DA2023001.

References

[1] B. G. Wybourne, Classical Groups for Physicists, Wiley Interscience, New York, 1974.

[2] A. Bohm, Y. Ne'eman, A.O. Barut, Dynamical Groups and Spectrum Generating Algebras, vols. 1 and 2, World Scientific, Singapore, 1988.

[3] M. J. Englefield, Group Theory and the Coulomb Problem, Wiley Interscience, USA, 1972.

[4] W. Miller, Lie Theory and Special Functions, Academic Press, New York, 1968.

[5] Y. Alhassid, J. Engel and J. Wu, Phys. Rev. Lett. 53(1984)17.

[6] Y. Alhassid, F. Iachello and R. D. Levine, Phys. Rev. Lett. 54(1985)1746.

[7] Y. Alhassid, J. Engel and F. Iachello, Phys. Rev. Lett. 57(1986)9.

[8] A. Frank and B. Wolf, Phys. Rev. Lett. 52(1984)1737.

[9] A. Frank, Y. Alhassid and F. Iachello L. Phys. Rev. A, 34(1986)677.

[10] A. Frank and B. Wolf, J. Math. Phys. 26(1985)973.

[11] J. Wu and Y. Alhassid, J. Math. Phys. 31(1986)557.

[12] G.A. Kerimov, Phys. Rev. Lett. 80(1998)2976.

[13] M. J. Englefield, J. Math. Phys. 28(1987)827.

[14] G. Lévai, J. Phys. A: Math. Gen. 27(1994)3809.

[15] D. Martínez, R.D. Mota, Ann. Phys. 323(2008)1024.

[16] D. Martínez, J.C. Flores-Urbina, R.D. Mota and V.D. Granados, J. Phys. A: Math. Theor. 43(2010)135201.

[17] J.D. Bjorken and S. T. Drell, Relativistic Quantum Mechanics, McGraw-Hill, New York, 1964.

[18] B. Thaller, The Dirac Equation, Springer-Verlag, Berlin, 1992.

[19] B. Thaller, Advanced Visual Quantum Mechanics, Springer-Verlag, Berlin, 2005.

[20] W. Greiner, Relativistic Quantum Mechanics, Springer-Verlag, Berlin, 1997.

[21] R.A. Swainson and G. W. F. Drake, J. Phys. A: Math. Gen. 24(1991)79.

[22] R.A. Swainson and G. W. F. Drake, J. Phys. A: Math. Gen. 24(1991)95.

[23] L. Infeld and T.E. Hull, Rev. Mod. Phys. 23(1951).

[24] J. Su, Phys. Rev. A 32(1985)3251.

[25] M.K.F. Wong, Phys. Rev. A 34(1986)1559.

[26] R. de Lima Rodrigues, Phys. Lett. A 326(2004)42.

[27] C.V. Sukumar, J. Phys. A: Math. Gen. 18(1985)L697.

[28] P.D. Jarvis and G.E. Stedman, J. Phys. A: Math. Gen. 19(1986)1373

[29] J.P. Dahl and T. Jorgensen, Int. J. Quant. Chem., 53(161)1995

[30] R.P. Martínez-y-Romero, J. Saldaña-Vega and A.L. Salas-Brito, J. Phys. A: Math. Gen. 31(1998)L157.

[31] R.P. Martínez-y-Romero and A.L. Salas-Brito and J. Saldaña-Vega, J. Math. Phys. 40(1999)2324.

[32] R.P. Martínez-y-Romero, H.N. Núñez-Yépez and A.L. Salas-Brito, Phys. Lett. A 339(2005)259.

[33] E. Schrödinger, *Proc. R. Ir. Acad. A* 46, 9(1940).

[34] D. Martínez, R.D. Mota, Ann. Phys. 323(2008)1024.

[35] B.G. Adams, J. Cizek and J. Paldus, Adv. Quant. Chem. Vol. 19 (1987)1. Reprinted in A. Bohm, Y. Neeman, A. O. Barut, Dynamical Groups and Spectrum Generating Algebras, vols. 1 and 2, World Scientific, Singapore, 1988.

[36] B.G. Adams, Algebraic Approach to Simple Quantum Systems, Springer-Verlag, Berlin, 1994.